


Lecture 13:

Camera Projection II


Reading: T&V Section 2.4

Recall: Imaging Geometry

**Object of Interest
in World Coordinate
System (U,V,W)**




Imaging Geometry



Camera Coordinate System (X,Y,Z).


- Z is optic axis
- Image plane located f units out along optic axis
- f is called focal length

Imaging Geometry



**Forward Projection onto image plane.
3D (X, Y, Z) projected to 2D (x, y)**

Imaging Geometry




Imaging Geometry


**Camera
Coordinates**

**Image (film)
Coordinates**

**World
Coordinates**



Forward Projection




We want a mathematical model to describe how 3D World points get projected into 2D Pixel coordinates.

Our goal: describe this sequence of transformations by a big matrix equation!

Intrinsic Camera Parameters


**World
Coords**

**U
V
W**




**Camera
Coords**

**X
Y
Z**



**Film
Coords**

*x
y*



**Pixel
Coords**

*u
v*

Affine Transformation


Intrinsic parameters

- Describes coordinate transformation between film coordinates (projected image) and pixel array
- Film cameras: scanning/digitization
- CCD cameras: grid of photosensors


still in T&V section 2.4

Intrinsic parameters (offsets)

film plane
(projected image)



pixel array




$$u = f \frac{X}{Z} + o_x \quad v = f \frac{Y}{Z} + o_y$$

o_x and o_y called image center or principle point

Intrinsic parameters


sometimes one or more coordinate axes are flipped (e.g. T&V section 2.4)



$$u = -f \frac{X}{Z} + o_x \quad v = -f \frac{Y}{Z} + o_y$$

Intrinsic parameters (scales)

sampling determines how many rows/cols in the image



Effective Scales: s_x and s_y

$$u = \frac{1}{s_x} f \frac{X}{Z} + o_x \quad v = \frac{1}{s_y} f \frac{Y}{Z} + o_y$$

Note, since we have different scale factors in x and y, we don't necessarily have square pixels!

Aspect ratio is s_y / s_x

Perspective projection matrix

Adding the intrinsic parameters into the perspective projection matrix:

$$\begin{bmatrix} x' \\ y' \\ z' \end{bmatrix} = \begin{bmatrix} f / s_x & 0 & o_x & 0 \\ 0 & f / s_y & o_y & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} X \\ Y \\ Z \\ 1 \end{bmatrix}$$

To verify:

$$\begin{aligned} u &= \frac{x'}{z'} \\ v &= \frac{y'}{z'} \end{aligned} \quad \Rightarrow \quad \begin{aligned} u &= \frac{1}{s_x} f \frac{X}{Z} + o_x \\ v &= \frac{1}{s_y} f \frac{Y}{Z} + o_y \end{aligned}$$

Note:

Sometimes, the image and the camera coordinate systems have opposite orientations: [the book does it this way]

$$\begin{aligned} f \frac{X}{Z} &= \downarrow (u - o_x) s_x \\ f \frac{Y}{Z} &= \downarrow (v - o_y) s_y \end{aligned} \quad \longrightarrow \quad \begin{bmatrix} x' \\ y' \\ z' \end{bmatrix} = \begin{bmatrix} -f/s_x & 0 & +o_x & 0 \\ 0 & -f/s_y & +o_y & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} X \\ Y \\ Z \\ 1 \end{bmatrix}$$

Note 2

In general, I like to think of the conversion as a separate 2D affine transformation from film coords (x,y) to pixel coordinates (u,v) :

$$\begin{pmatrix} u' \\ v' \\ w' \end{pmatrix} = \underbrace{\begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ 0 & 0 & 1 \end{pmatrix}}_{\mathbf{M}_{\text{aff}}} \underbrace{\begin{bmatrix} f & 0 & 0 & 0 \\ 0 & f & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}}_{\mathbf{M}_{\text{proj}}} \begin{bmatrix} X \\ Y \\ Z \\ 1 \end{bmatrix}$$

$$\mathbf{u} = \mathbf{M}_{\text{int}} \mathbf{P}_C = \mathbf{M}_{\text{aff}} \mathbf{M}_{\text{proj}} \mathbf{P}_C$$




Huh?

Did he just say it was “a fine” transformation?

No, it was “affine” transformation, a type of 2D to 2D mapping defined by 6 parameters.

More on this in a moment...

Summary : Forward Projection



Lecture 13/14:

Intro to Image Mappings

Image Mappings Overview





FIGURE 1. Basic set of 2D planar transformations


Geometric Image Mappings



$$\begin{aligned}x' &= f(x, y, \{\text{parameters}\}) \\y' &= g(x, y, \{\text{parameters}\})\end{aligned}$$

Linear Transformations


(Can be written as matrices)



$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} & & \\ & & \\ & & \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

$\mathbf{M}(\text{params})$

Translation




$$\begin{aligned}x' &= x + t_x \\y' &= y + t_y\end{aligned}$$

equations

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

matrix form

Scale




$$\begin{aligned}x'_i &= s x_i \\y'_i &= s y_i\end{aligned}$$

equations

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} s & 0 & 0 \\ 0 & s & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

matrix form

Rotation




$$\begin{aligned}x' &= x_i \cos \theta - y_i \sin \theta \\y' &= x_i \sin \theta + y_i \cos \theta\end{aligned}$$

equations

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

matrix form

Euclidean (Rigid)



$$\begin{aligned}x' &= x_i \cos \theta - y_i \sin \theta + t_x \\y' &= x_i \sin \theta + y_i \cos \theta + t_y\end{aligned}$$

equations

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} \cos \theta & -\sin \theta & t_x \\ \sin \theta & \cos \theta & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

matrix form

Partitioned Matrices

A *partitioned matrix*, or a *block matrix*, is a *matrix* M that has been constructed from other smaller matrices. These smaller matrices are called *blocks* or *sub-matrices* of M .

For instance, if we *partition* the below 5×5 matrix as follows

$$L = \left(\begin{array}{cc|ccc} 1 & 0 & 1 & 2 & 3 \\ 0 & 1 & 1 & 2 & 3 \\ \hline 2 & 3 & 9 & 9 & 9 \\ 2 & 3 & 9 & 9 & 9 \\ 2 & 3 & 9 & 9 & 9 \end{array} \right),$$

then we can define the matrices

$$A = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, B = \begin{pmatrix} 1 & 2 & 3 \\ 1 & 2 & 3 \end{pmatrix}, C = \begin{pmatrix} 2 & 3 \\ 2 & 3 \\ 2 & 3 \end{pmatrix}, D = \begin{pmatrix} 9 & 9 & 9 \\ 9 & 9 & 9 \\ 9 & 9 & 9 \end{pmatrix}$$

and write L as

$$L = \begin{pmatrix} A & B \\ C & D \end{pmatrix}, \text{ or } L = \left(\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right).$$

Partitioned Matrices

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \left[\begin{array}{cc|c} \cos \theta & -\sin \theta & t_x \\ \sin \theta & \cos \theta & t_y \\ \hline 0 & 0 & 1 \end{array} \right] \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} \overset{2 \times 1}{p'} \\ \underset{1 \times 1}{1} \end{bmatrix} = \begin{bmatrix} \overset{2 \times 2}{R} & \overset{2 \times 1}{t} \\ \underset{1 \times 2}{0} & \underset{1 \times 1}{1} \end{bmatrix} \begin{bmatrix} \overset{2 \times 1}{p} \\ \underset{1 \times 1}{1} \end{bmatrix}$$

matrix form

$$p' = Rp + t$$

equation form


Another Example (from last time)

$$\begin{pmatrix} X \\ Y \\ Z \\ \hline 1 \end{pmatrix} = \begin{pmatrix} r_{11} & r_{12} & r_{13} & | & t_x \\ r_{21} & r_{22} & r_{23} & | & t_y \\ r_{31} & r_{32} & r_{33} & | & t_z \\ \hline 0 & 0 & 0 & | & 1 \end{pmatrix} \begin{pmatrix} U \\ V \\ W \\ \hline 1 \end{pmatrix}$$

$$\begin{pmatrix} \overset{3 \times 1}{\mathbf{P}_C} \\ \overset{1 \times 1}{1} \end{pmatrix} = \begin{pmatrix} \overset{3 \times 3}{\mathbf{R}} & \overset{3 \times 1}{\mathbf{T}} \\ \overset{1 \times 3}{0} & \overset{1 \times 1}{1} \end{pmatrix} \begin{pmatrix} \overset{3 \times 1}{\mathbf{P}_W} \\ \overset{1 \times 1}{1} \end{pmatrix}$$

$$\mathbf{P}_C = \mathbf{R} \mathbf{P}_W + \mathbf{T}$$

Similarity (scaled Euclidean)




$$p' = sRp + t$$

equations

$$\begin{bmatrix} p' \\ 1 \end{bmatrix} = \begin{bmatrix} sR & t \\ 0 & 1 \end{bmatrix} \begin{bmatrix} p \\ 1 \end{bmatrix}$$

matrix form

Affine




$$p' = Ap + b$$

equations



$$\begin{bmatrix} p' \\ 1 \end{bmatrix} = \begin{bmatrix} A & b \\ 0 & 1 \end{bmatrix} \begin{bmatrix} p \\ 1 \end{bmatrix}$$

matrix form

Projective



transform



Note!


$$p' = \frac{Ap + b}{c^T p + 1}$$

equations


$$\begin{bmatrix} p' \\ 1 \end{bmatrix} \sim \begin{bmatrix} A & b \\ c^T & 1 \end{bmatrix} \begin{bmatrix} p \\ 1 \end{bmatrix}$$

matrix form


Summary of 2D Transformations




rotation




translation




scale



aspect ratio







skew



perspective
warp





Summary of 2D Transformations

Euclidean




Summary of 2D Transformations

Similarity




Summary of 2D Transformations


Affine




rotation




translation




scale



aspect ratio




skew







perspective
warp

Summary of 2D Transformations

Projective



Summary of 2D Transformations

Name	Matrix	# D.O.F.	Preserves:	Icon
translation	$[\mathbf{I} \mid \mathbf{t}]_{2 \times 3}$	2	orientation + ...	
rigid (Euclidean)	$[\mathbf{R} \mid \mathbf{t}]_{2 \times 3}$	3	lengths + ...	
similarity	$[s\mathbf{R} \mid \mathbf{t}]_{2 \times 3}$	4	angles + ...	
affine	$[\mathbf{A}]_{2 \times 3}$	6	parallelism + ...	
projective	$[\mathbf{H}]_{3 \times 3}$	8	straight lines	